arXiv:0706.4160v1 [math.DG] 28 Jun 2007

EXPLICIT FORMULAS FOR BIHARMONIC SUBMANIFOLDS IN SASAKIAN SPACE FORMS

D. FETCU AND C. ONICIUC

Dedicated to Professor Neculai Papaghiuc on his 60-th birthday

Abstract. We classify the biharmonic Legendre curves in a Sasakian space form, and obtain their explicit parametric equations in the $(2n+1)$-dimensional unit sphere endowed with the canonical and deformed Sasakian structures defined by Tanno. Then, composing with the flow of the Reeb vector field, we transform a biharmonic integral submanifold into a biharmonic anti-invariant submanifold. Using this method we obtain new examples of biharmonic submanifolds in spheres and, in particular, in $\mathbb{S}^7$.

1. Introduction

Biharmonic maps between Riemannian manifolds $\phi : (M,g) \to (N,h)$ are the critical points of the *bienergy functional* $E_2(\phi) = \frac{1}{2}\int_M |\tau(\phi)|^2 \ v_g$. They represent a natural generalization of the well-known *harmonic maps* ([8]), the critical points of the *energy functional* $E(\phi) = \frac{1}{2}\int_M |d\phi|^2 \ v_g$, and of the biharmonic submanifolds in Euclidean spaces defined by B.-Y. Chen ([7]).

The Euler-Lagrange equation for the energy functional is $\tau(\phi) = 0$, where $\tau(\phi) = \operatorname{trace} \nabla d\phi$ is the tension field, and the Euler-Lagrange equation for the bienergy functional was derived by G. Y. Jiang in [14]:

$$\begin{aligned} \tau_2(\phi) &= -\Delta\tau(\phi) - \operatorname{trace} \ R^N(d\phi, \tau(\phi))d\phi \\ &= 0. \end{aligned}$$

Since any harmonic map is biharmonic, we are interested in non-harmonic biharmonic maps, which are called *proper-biharmonic*.

A special case of biharmonic maps is represented by the biharmonic Riemannian immersions, or biharmonic submanifolds. There are several results of classification or construction for such submanifolds in space forms ([15], [5]). Then, the next step would be the study of biharmonic submanifolds in Sasakian space forms. In this context J. Inoguchi classified in [13] the proper-biharmonic Legendre curves and Hopf cylinders in a 3-dimensional Sasakian space form $M^3(c)$, and in [11] the explicit parametric equations were obtained. Then, T. Sasahara and his collaborators studied the biharmonic integral surfaces and 3-dimensional biharmonic anti-invariant submanifolds in $\mathbb{S}^5$ ([1], [17]).

Recent results on biharmonic submanifolds in spaces of nonconstant sectional curvature were obtained by T. Ichiyama, J. Inoguchi and H. Urakawa in [12], by Y.-L. Ou and Z.-P. Wang in [16], and by W. Zhang in [20].

2000 *Mathematics Subject Classification.* 53C42, 53B25.

Key words and phrases. Biharmonic submanifolds, Sasakian space forms, Legendre curves, integral submanifolds.

The authors were partially supported by the Grant CEEX, ET, 5871/2006, Romania.

Biharmonic submanifolds in pseudo-Euclidean spaces were also studied, and many examples and classification results were obtained (for example, see [2], [7]).

The goals of our paper are to obtain new classification results for biharmonic Legendre curves in any dimensional Sasakian space form and to provide a method for constructing biharmonic submanifolds. In order to obtain explicit examples, we use the $(2n+1)$-dimensional unit sphere $\mathbb{S}^{2n+1}$ as a model of Sasakian space form.

For a general account of biharmonic maps see [15] and *The Bibliography of Biharmonic Maps* [18].

Conventions. We work in the C^∞ category, that means manifolds, metrics, connections and maps are smooth. The Lie algebra of the vector fields on M is denoted by $C(TM)$.

2. PRELIMINARIES

2.1. **Contact manifolds.** A *contact metric structure* on a manifold N^{2n+1} is given by (φ, ξ, η, g), where φ is a tensor field of type $(1,1)$ on N, ξ is a vector field, η is an 1-form and g is a Riemannian metric such that

$$\varphi^2 = -I + \eta \otimes \xi, \quad \eta(\xi) = 1,$$

$$g(\varphi X, \varphi Y) = g(X,Y) - \eta(X)\eta(Y), \quad g(X, \varphi Y) = d\eta(X,Y), \quad \forall X, Y \in C(TN).$$

A contact metric structure (φ, ξ, η, g) is called *normal* if

$$N_\varphi + 2d\eta \otimes \xi = 0,$$

where

$$N_\varphi(X,Y) = [\varphi X, \varphi Y] - \varphi[\varphi X, Y] - \varphi[X, \varphi Y] + \varphi^2[X,Y], \quad \forall X, Y \in C(TN),$$

is the Nijenhuis tensor field of φ.

A contact metric manifold $(N, \varphi, \xi, \eta, g)$ is a *Sasakian manifold* if it is normal or, equivalently, if

$$(\nabla_X \varphi)(Y) = g(X,Y)\xi - \eta(Y)X, \quad \forall X, Y \in C(TN).$$

The *contact distribution* of a Sasakian manifold $(N, \varphi, \xi, \eta, g)$ is defined by $\{X \in TN : \eta(X) = 0\}$, and an integral curve of the contact distribution is called *Legendre curve*. A submanifold M of N which is tangent to ξ is said to be *anti-invariant* if φ maps any vector tangent to M and normal to ξ to a vector normal to M.

Let $(N, \varphi, \xi, \eta, g)$ be a Sasakian manifold. The sectional curvature of a 2-plane generated by X and φX, where X is an unit vector orthogonal to ξ, is called *φ-sectional curvature* determined by X. A Sasakian manifold with constant φ-sectional curvature c is called a *Sasakian space form* and it is denoted by $N(c)$.

The curvature tensor field of a Sasakian space form $N(c)$ is given by

$$\begin{aligned} R(X,Y)Z = \ & \tfrac{c+3}{4}\{g(Z,Y)X - g(Z,X)Y\} + \tfrac{c-1}{4}\{\eta(Z)\eta(X)Y - \\ & -\eta(Z)\eta(Y)X + g(Z,X)\eta(Y)\xi - g(Z,Y)\eta(X)\xi + \\ & +g(Z,\varphi Y)\varphi X - g(Z,\varphi X)\varphi Y + 2g(X,\varphi Y)\varphi Z\}. \end{aligned} \tag{2.1}$$

Let $\mathbb{S}^{2n+1} = \{z \in \mathbb{C}^{n+1} : |z| = 1\}$ be the unit $(2n+1)$-dimensional sphere endowed with its standard metric field g_0. Consider the following structure tensor fields on $\mathbb{S}^{2n+1}$: $\xi_0 = -\mathcal{I}z$ for each $z \in \mathbb{S}^{2n+1}$, where $\mathcal{I}$ is the usual almost complex structure on $\mathbb{C}^{n+1}$ defined by

$$\mathcal{I}z = (-y^1, ..., -y^{n+1}, x^1, ..., x^{n+1}),$$

for $z=(x^1,...,x^{n+1},y^1,...,y^{n+1})$, and $\varphi_0=s\circ\mathcal{I}$, where $s:T_z\mathbb{C}^{n+1}\to T_z\mathbb{S}^{2n+1}$ denotes the orthogonal projection. Equipped with these tensors, $\mathbb{S}^{2n+1}$ becomes a Sasakian space form with φ_0-sectional curvature equal to 1 ([3]).
Now, consider the deformed structure on $\mathbb{S}^{2n+1}$, introduced by Tanno in [13],

$$\eta=a\eta_0,\quad \xi=\frac{1}{a}\xi_0,\quad \varphi=\varphi_0,\quad g=ag_0+a(a-1)\eta_0\otimes\eta_0,$$

where a is a positive constant. The structure (φ,ξ,η,g) is still a Sasakian structure and $(\mathbb{S}^{2n+1},\varphi,\xi,\eta,g)$ is a Sasakian space form with constant φ-sectional curvature $c=\frac{4}{a}-3$ ([1]).

2.2. **3-Sasakian manifolds.** If a manifold N admits three Sasakian structures $(\varphi_a,\xi_a,\eta_a,g)$, $a=1,2,3$, satisfying

$$\varphi_c=-\varphi_a\varphi_b+\eta_b\otimes\xi_a=\varphi_b\varphi_a-\eta_a\otimes\xi_b,$$

$$\xi_c=-\varphi_a\xi_b=\varphi_b\xi_a,\quad \eta_c=-\eta_a\circ\varphi_b=\eta_b\circ\varphi_a,$$

for an even permutation (a,b,c) of $(1,2,3)$, then the manifold is said to have a *Sasakian 3-structure*. The dimension of such a manifold is of the form $4n+3$. We note that the maximum dimension of a submanifold of a 3-Sasakian manifold N^{4n+3} which is an integral submanifold with respect to all three Sasakian structures is n.

3. Biharmonic Legendre curves in Sasakian space forms

Definition 3.1. Let (N^m,g) be a Riemannian manifold and $\gamma:I\to N$ a curve parametrized by arc length, that is $|\gamma'|=1$. Then γ is called a *Frenet curve of osculating order* r, $1\le r\le m$, if there exists orthonormal vector fields $E_1,E_2,...,E_r$ along γ such that

$$\begin{cases} E_1=\gamma'=T\\ \nabla_TE_1=\kappa_1E_2\\ \nabla_TE_2=-\kappa_1E_1+\kappa_2E_3\\ \dots\\ \nabla_TE_r=-\kappa_{r-1}E_{r-1}\end{cases},$$

where $\kappa_1,...,\kappa_{r-1}$ are positive functions on I.

Remark 3.2. A geodesic is a Frenet curve of osculating order 1; a *circle* is a Frenet curve of osculating order 2 with $\kappa_1=$ constant; a *helix of order* r, $r\ge3$, is a Frenet curve of osculating order r with $\kappa_1,...,\kappa_{r-1}$ constants; a helix of order 3 is called, simply, helix.

Now let $(N^{2n+1},\varphi,\xi,\eta,g)$ be a Sasakian space form with constant φ-sectional curvature c and $\gamma:I\to N$ a Legendre Frenet curve of osculating order r. As

$$\begin{aligned}\nabla^3_TT=\ &(-3\kappa_1\kappa_1')E_1+(\kappa_1''-\kappa_1^3-\kappa_1\kappa_2^2)E_2+(2\kappa_1'\kappa_2+\kappa_1\kappa_2')E_3\\ &+\kappa_1\kappa_2\kappa_3E_4\end{aligned}\tag{3.1}$$

and

$$R(T,\nabla_TT)T=-\frac{(c+3)\kappa_1}{4}E_2-\frac{3(c-1)\kappa_1}{4}g(E_2,\varphi T)\varphi T,\tag{3.2}$$

we get

$$\begin{aligned}\tau_2(\gamma) = {} & \nabla_T^3 T - R(T,\nabla_T T)T \\ = {} & (-3\kappa_1\kappa_1')E_1 + \Big(\kappa_1'' - \kappa_1^3 - \kappa_1\kappa_2^2 + \tfrac{(c+3)\kappa_1}{4}\Big)E_2 \\ & +(2\kappa_1'\kappa_2 + \kappa_1\kappa_2')E_3 + \kappa_1\kappa_2\kappa_3 E_4 + \tfrac{3(c-1)\kappa_1}{4} g(E_2,\varphi T)\varphi T.\end{aligned} \tag{3.3}$$

In the following we shall solve the biharmonic equation $\tau_2(\gamma) = 0$. The problem is to find the relation between φT and the Frenet frame field. The simplest two cases are provided by $\frac{3(c-1)\kappa_1}{4} g(E_2,\varphi T) = 0$. So,

Case I: c = 1.

In this case γ is proper-biharmonic if and only if

$$\begin{cases} \kappa_1 = \text{constant} > 0, \quad \kappa_2 = \text{constant} \\ \kappa_1^2 + \kappa_2^2 = 1 \\ \kappa_2\kappa_3 = 0 \end{cases}.$$

One obtains

Theorem 3.3. *If $c = 1$ and $n \geq 2$, then γ is proper-biharmonic if and only if either γ is a circle with $\kappa_1 = 1$, or γ is a helix with $\kappa_1^2 + \kappa_2^2 = 1$.*

Remark 3.4. If $n = 1$ and γ is a non-geodesic Legendre curve we have $\nabla_T T = \pm\kappa_1\varphi T$ and then $E_2 = \pm\varphi T$ and $\nabla_T E_2 = \pm\nabla_T\varphi T = \pm(\xi \mp \kappa_1 T) = -\kappa_1 T \pm \xi$. Therefore $\kappa_2 = 1$ and γ cannot be biharmonic.

Case II: $\mathbf{c \neq 1}$, $\mathbf{E_2 \perp \varphi T}$.

In this case γ is proper-biharmonic if and only if

$$\begin{cases} \kappa_1 = \text{constant} > 0, \quad \kappa_2 = \text{constant} \\ \kappa_1^2 + \kappa_2^2 = \frac{c+3}{4} \\ \kappa_2\kappa_3 = 0 \end{cases}.$$

Before stating the theorem we need the following

Lemma 3.5. *Let γ be a Legendre Frenet curve of osculating order 3 and $E_2 \perp \varphi T$. Then $\{T = E_1, E_2, E_3, \varphi T, \xi, \nabla_T\varphi T\}$ are linearly independent, in any point, and hence $n \geq 3$.*

Proof. Since γ is a Frenet curve of osculating order 3, we have

$$\begin{cases} E_1 = \gamma' = T \\ \nabla_T E_1 = \kappa_1 E_2 \\ \nabla_T E_2 = -\kappa_1 E_1 + \kappa_2 E_3 \\ \nabla_T E_3 = -\kappa_2 E_2 \end{cases}.$$

It is easy to see that, in an arbitrary point, the system

$$S_1 = \{T = E_1, E_2, E_3, \varphi T, \xi, \nabla_T\varphi T\}$$

has only non-zero vectors and

$$T \perp E_2, \ \ T \perp E_3, \ \ T \perp \varphi T, \ \ T \perp \xi, \ \ T \perp \nabla_T \varphi T.$$

Thus S_1 is linearly independent if and only if $S_2 = \{E_2, E_3, \varphi T, \xi, \nabla_T \varphi T\}$ is linearly independent. Further, as

$$E_2 \perp \xi, \ E_2 \perp \nabla_T \varphi T, \ E_3 \perp \xi, \ E_3 \perp \nabla_T \varphi T, \ \varphi T \perp \xi, \ \varphi T \perp \nabla_T \varphi T,$$

and

$$E_2 \perp E_3 \perp \varphi T,$$

it follows that S_2 is linearly independent if and only if $S_3 = \{\xi, \nabla_T \varphi T\}$ is linearly independent. But $\nabla_T \varphi T = \xi + \kappa_1 \varphi E_2$, $\kappa_1 \neq 0$, and therefore S_3 is linearly independent. □

Now we can state

Theorem 3.6. *Assume that $c \neq 1$ and $\nabla_T T \perp \varphi T$. We have*
1) If $c \leq -3$ then γ is biharmonic if and only if it is a geodesic.
2) If $c > -3$ then γ is proper-biharmonic if and only if either
a) $n \geq 2$ and γ is a circle with $\kappa_1^2 = \frac{c+3}{4}$. In this case $\{E_1, E_2, \varphi T, \xi\}$ are linearly independent,
or
b) $n \geq 3$ and γ is a helix with $\kappa_1^2 + \kappa_2^2 = \frac{c+3}{4}$. In this case $\{E_1, E_2, E_3, \varphi T, \xi, \nabla_T \varphi T\}$ are linearly independent.

Case III: $\mathbf{c \neq 1}$, $\mathbf{E_2 \parallel \varphi T}$.
In this case γ is proper-biharmonic if and only if

$$\begin{cases} \kappa_1 = \text{constant} > 0, \quad \kappa_2 = \text{constant} \\ \\ \kappa_1^2 + \kappa_2^2 = c \\ \\ \kappa_2 \kappa_3 = 0 \end{cases}.$$

We can assume that $E_2 = \varphi T$. Then we have $\nabla_T T = \kappa_1 E_2 = \kappa_1 \varphi T$, $\nabla_T E_2 = \nabla_T \varphi T = \xi - \kappa_1 T$. That means $E_3 = \xi$ and $\kappa_2 = 1$. Hence $\nabla_T E_3 = \nabla_T \xi = -\varphi T = -E_2$.
Therefore

Theorem 3.7. *If $c \neq 1$ and $\nabla_T T \parallel \varphi T$, then $\{T, \varphi T, \xi\}$ is the Frenet frame field of γ and we have*
1) If $c \leq 1$ then γ is biharmonic if and only if it is a geodesic.
2) If $c > 1$ then γ is proper-biharmonic if and only if it is a helix with $\kappa_1^2 = c - 1$ (and $\kappa_2 = 1$).

Remark 3.8. If $n = 1$ then $\nabla_T T \parallel \varphi T$ and we reobtain Inoguchi's result [13].

Case IV: $\mathbf{c \neq 1}$ and $\mathbf{g(E_2, \varphi T)}$ is not constant $\mathbf{0, 1}$ or $\mathbf{-1}$.
Assume that γ is a Legendre Frenet curve of osculating order r, $4 \leq r \leq 2n+1$, $n \geq 2$. If γ is biharmonic it follows that $\varphi T \in \operatorname{span}\{E_2, E_3, E_4\}$.
Now, we denote $f(t) = g(E_2, \varphi T)$ and differentiating along γ we obtain

$$\begin{aligned} f'(t) &= g(\nabla_T E_2, \varphi T) + g(E_2, \nabla_T \varphi T) = g(\nabla_T E_2, \varphi T) + g(E_2, \xi + \kappa_1 \varphi E_2) \\ &= g(\nabla_T E_2, \varphi T) = g(-\kappa_1 T + \kappa_2 E_3, \varphi T) \\ &= \kappa_2 g(E_3, \varphi T). \end{aligned}$$

Since $\varphi T = g(\varphi T, E_2)E_2 + g(\varphi T, E_3)E_3 + g(\varphi T, E_4)E_4$, the curve γ is proper-biharmonic if and only if

$$\begin{cases} \kappa_1 = \text{constant} > 0 \\ \kappa_1^2 + \kappa_2^2 = \frac{c+3}{4} + \frac{3(c-1)}{4} f^2 \\ \kappa_2' = -\frac{3(c-1)}{4} f g(\varphi T, E_3) \\ \kappa_2 \kappa_3 = -\frac{3(c-1)}{4} f g(\varphi T, E_4) \end{cases}.$$

Using the expression of $f'(t)$ we see that the third equation of the above system is equivalent to

$$\kappa_2^2 = -\frac{3(c-1)}{4} f^2 + \omega_0,$$

where $\omega_0 = \text{constant}$. Replacing in the second equation it follows

$$\kappa_1^2 = \frac{c+3}{4} - \omega_0 + \frac{3(c-1)}{2} f^2,$$

which implies $f = \text{constant}$. Thus $\kappa_2 = \text{constant} > 0$, $g(E_3, \varphi T) = 0$ and then $\varphi T = f E_2 + g(\varphi T, E_4)E_4$. It follows that there exists an unique constant $\alpha_0 \in (0, 2\pi) \setminus \{\frac{\pi}{2}, \pi, \frac{3\pi}{2}\}$ such that $f = \cos \alpha_0$ and $g(\varphi T, E_4) = \sin \alpha_0$.
We can state

Theorem 3.9. *Let $c \neq 1$, $n \geq 2$ and γ a Legendre Frenet curve of osculating order $r \geq 4$ such that $g(E_2, \varphi T)$ is not constant $0, 1$ or -1. We have*

a) If $c \leq -3$ then γ is biharmonic if and only if it is a geodesic.

b) If $c > -3$ then γ is proper-biharmonic if and only if $\varphi T = \cos \alpha_0 E_2 + \sin \alpha_0 E_4$ and

$$\begin{cases} \kappa_1 = \text{constant} > 0, \quad \kappa_2 = \text{constant} \\ \kappa_1^2 + \kappa_2^2 = \frac{c+3}{4} + \frac{3(c-1)}{4} \cos^2 \alpha_0 \\ \kappa_2 \kappa_3 = -\frac{3(c-1)}{8} \sin 2\alpha_0 \end{cases},$$

where $\alpha_0 \in (0, 2\pi) \setminus \{\frac{\pi}{2}, \pi, \frac{3\pi}{2}\}$ is a constant such that $c + 3 + 3(c-1)\cos^2 \alpha_0 > 0$ and $3(c-1)\sin 2\alpha_0 < 0$.

Remark 3.10. In this case we may obtain biharmonic curves which are not helices.

Remark 3.11. We note that a preliminary version of the full classification of the proper-biharmonic Legendre curves in Sasakian space forms was obtained in [10].

In the following, we shall choose the unit $(2n+1)$-dimensional sphere $\mathbb{S}^{2n+1}$ with its canonical and modified Sasakian structures as a model for the complete, simply connected Sasakian space form with constant φ-sectional curvature $c > -3$, and we will find the explicit equations of biharmonic Legendre curves obtained in the first three cases, viewed as curves in $\mathbb{R}^{2n+2}$.

Theorem 3.12. *Let $\gamma : I \to (\mathbb{S}^{2n+1}, \varphi_0, \xi_0, \eta_0, g_0)$, $n \geq 2$, be a proper-biharmonic Legendre curve parametrized by arc length. Then the equation of γ in the Euclidean space $\mathbb{E}^{2n+2} = (\mathbb{R}^{2n+2}, \langle , \rangle)$, is either*

$$\gamma(s) = \frac{1}{\sqrt{2}} \cos\left(\sqrt{2}s\right) e_1 + \frac{1}{\sqrt{2}} \sin\left(\sqrt{2}s\right) e_2 + \frac{1}{\sqrt{2}} e_3$$

where $\{e_i, \mathcal{I}e_j\}$ *are constant unit vectors orthogonal to each other, or*

$$\gamma(s) = \frac{1}{\sqrt{2}}\cos(As)e_1 + \frac{1}{\sqrt{2}}\sin(As)e_2 + \frac{1}{\sqrt{2}}\cos(Bs)e_3 + \frac{1}{\sqrt{2}}\sin(Bs)e_4,$$

where

$$A = \sqrt{1+\kappa_1}, \quad B = \sqrt{1-\kappa_1}, \quad \kappa_1 \in (0,1), \tag{3.4}$$

and $\{e_i\}$ *are constant unit vectors orthogonal to each other, with*

$$\langle e_1, \mathcal{I}e_3\rangle = \langle e_1, \mathcal{I}e_4\rangle = \langle e_2, \mathcal{I}e_3\rangle = \langle e_2, \mathcal{I}e_4\rangle = 0, \quad A\langle e_1, \mathcal{I}e_2\rangle + B\langle e_3, \mathcal{I}e_4\rangle = 0.$$

Proof. Let us denote by $\dot{\nabla}$ and by $\widetilde{\nabla}$ the Levi-Civita connections on $(\mathbb{S}^{2n+1}, g_0)$ and $(\mathbb{R}^{2n+2}, \langle,\rangle)$, respectively.

First, assume that γ is the biharmonic circle, that is $\kappa_1 = 1$. From the Gauss and Frenet equations we get

$$\widetilde{\nabla}_T T = \dot{\nabla}_T T - \langle T, T\rangle\gamma = \kappa_1 E_2 - \gamma$$

and

$$\widetilde{\nabla}_T \widetilde{\nabla}_T T = (-\kappa_1^2 - 1)T = -2T,$$

which implies

$$\gamma''' + 2\gamma' = 0.$$

The general solution of the above equation is

$$\gamma(s) = \cos\left(\sqrt{2}s\right)c_1 + \sin\left(\sqrt{2}s\right)c_2 + c_3,$$

where $\{c_i\}$ are constant vectors in $\mathbb{E}^{2n+2}$.
Now, as γ satisfies

$$\langle\gamma,\gamma\rangle = 1,\ \langle\gamma',\gamma'\rangle = 1,\ \langle\gamma,\gamma'\rangle = 0,\ \langle\gamma',\gamma''\rangle = 0,\ \langle\gamma'',\gamma''\rangle = 2,\ \langle\gamma,\gamma''\rangle = -1,$$

and since in $s = 0$ we have $\gamma = c_1 + c_3$, $\gamma' = \sqrt{2}c_2$, $\gamma'' = -2c_1$, we obtain

$$c_{11} + 2c_{13} + c_{33} = 1,\ c_{22} = \frac{1}{2},\ c_{12} + c_{23} = 0,\ c_{12} = 0,\ c_{11} = \frac{1}{2}, c_{11} + c_{13} = \frac{1}{2},$$

where $c_{ij} = \langle c_i, c_j\rangle$. Further, we get that $\{c_i\}$ are orthogonal vectors in $\mathbb{E}^{2n+2}$ with $|c_1| = |c_2| = |c_3| = \frac{1}{\sqrt{2}}$.
Finally, using the fact that γ is a Legendre curve one obtains easily that $\langle c_i, \mathcal{I}c_j\rangle = 0$ for any $i, j = 1, 2, 3$.

Suppose now γ is the biharmonic helix, that is $\kappa_1^2 + \kappa_2^2 = 1$, $\kappa_1 \in (0,1)$. From the Gauss and Frenet equations we get

$$\widetilde{\nabla}_T T = \dot{\nabla}_T T - \langle T, T\rangle\gamma = \kappa_1 E_2 - \gamma,$$

$$\widetilde{\nabla}_T \widetilde{\nabla}_T T = \kappa_1 \widetilde{\nabla}_T E_2 - T = \kappa_1\Big(-\kappa_1 T + \kappa_2 E_3\Big) - T = -\Big(\kappa_1^2 + 1\Big)T + \kappa_1\kappa_2 E_3$$

and

$$\widetilde{\nabla}_T \widetilde{\nabla}_T \widetilde{\nabla}_T T = -\Big(\kappa_1^2+1\Big)\widetilde{\nabla}_T T + \kappa_1\kappa_2\widetilde{\nabla}_T E_3 = -\Big(\kappa_1^2+1\Big)\widetilde{\nabla}_T T - \kappa_1\kappa_2^2 E_2 = -2\gamma'' - \kappa_2^2\gamma.$$

Hence

$$\gamma^{iv} + 2\gamma'' + \kappa_2^2\gamma = 0,$$

and its general solution is

$$\gamma(s) = \cos(As)c_1 + \sin(As)c_2 + \cos(Bs)c_3 + \sin(Bs)c_4,$$

where A, B are given by (3.4) and $\{c_i\}$ are constant vectors in $\mathbb{E}^{2n+2}$.

As γ satisfies

$$\langle\gamma,\gamma\rangle=1,\ \langle\gamma',\gamma'\rangle=1,\ \langle\gamma,\gamma'\rangle=0,\ \langle\gamma',\gamma''\rangle=0,\ \langle\gamma'',\gamma''\rangle=1+\kappa_1^2,$$

$$\langle\gamma,\gamma''\rangle=-1,\ \langle\gamma',\gamma'''\rangle=-(1+\kappa_1^2),\ \langle\gamma'',\gamma'''\rangle=0,\ \langle\gamma,\gamma'''\rangle=0,\ \langle\gamma''',\gamma'''\rangle=3\kappa_1^2+1,$$

and since in $s=0$ we have $\gamma=c_1+c_3$, $\gamma'=Ac_2+Bc_4$, $\gamma''=-A^2c_1-B^2c_3$, $\gamma'''=-A^3c_2-B^3c_4$, we obtain

$$c_{11}+2c_{13}+c_{33}=1 \tag{3.5}$$

$$A^2c_{22}+2ABc_{24}+B^2c_{44}=1 \tag{3.6}$$

$$Ac_{12}+Ac_{23}+Bc_{14}+Bc_{34}=0 \tag{3.7}$$

$$A^3c_{12}+AB^2c_{23}+A^2Bc_{14}+B^3c_{34}=0 \tag{3.8}$$

$$A^4c_{11}+2A^2B^2c_{13}+B^4c_{33}=1+\kappa_1^2 \tag{3.9}$$

$$A^2c_{11}+(A^2+B^2)c_{13}+B^2c_{33}=1 \tag{3.10}$$

$$A^4c_{22}+(AB^3+A^3B)c_{24}+B^4c_{44}=1+\kappa_1^2 \tag{3.11}$$

$$A^5c_{12}+A^3B^2c_{23}+A^2B^3c_{14}+B^5c_{34}=0 \tag{3.12}$$

$$A^3c_{12}+A^3c_{23}+B^3c_{14}+B^3c_{34}=0 \tag{3.13}$$

$$A^6c_{22}+2A^3B^3c_{24}+B^6c_{44}=3\kappa_1^2+1 \tag{3.14}$$

where $c_{ij}=\langle c_i,c_j\rangle$. Since the determinant of the system given by (3.7), (3.8), (3.12) and (3.13) is $-A^2B^2(A^2-B^2)^4\neq 0$ it follows that

$$c_{12}=c_{23}=c_{14}=c_{34}=0.$$

The equations (3.5), (3.9) and (3.10) give

$$c_{11}=\frac{1}{2},\quad c_{13}=0,\quad c_{33}=\frac{1}{2},$$

and, from (3.6), (3.11) and (3.14) follows that

$$c_{22}=\frac{1}{2},\quad c_{24}=0,\quad c_{44}=\frac{1}{2}.$$

Therefore, we obtain that $\{c_i\}$ are orthogonal vectors in $\mathbb{E}^8$ with $|c_1|=|c_2|=|c_3|=|c_4|=\frac{1}{\sqrt{2}}$.

Finally, since γ is a Legendre curve one obtains the conclusion of the Theorem. □

Theorem 3.13. *Let* $\gamma: I\to(\mathbb{S}^{2n+1},\varphi,\xi,\eta,g)$, $n\geq 2$, $a>0$, $a\neq 1$, *be a proper-biharmonic Legendre curve parametrized by arc length such that* $g(\nabla_{\gamma'}\gamma',\varphi\gamma')=0$. *Then the equation of* γ *in the Euclidean space* $\mathbb{E}^{2n+2}$, *is either*

$$\gamma(s)=\frac{1}{\sqrt{2}}\cos\Big(\sqrt{\frac{2}{a}}s\Big)e_1+\frac{1}{\sqrt{2}}\sin\Big(\sqrt{\frac{2}{a}}s\Big)e_2+\frac{1}{\sqrt{2}}e_3$$

for $n\geq 2$ *or, for* $n\geq 3$,

$$\gamma(s)=\frac{1}{\sqrt{2}}\cos(As)e_1+\frac{1}{\sqrt{2}}\sin(As)e_2+\frac{1}{\sqrt{2}}\cos(Bs)e_3+\frac{1}{\sqrt{2}}\sin(Bs)e_4,$$

where

$$A=\sqrt{\frac{1+\kappa_1\sqrt{a}}{a}},\quad B=\sqrt{\frac{1-\kappa_1\sqrt{a}}{a}},\quad \kappa_1\in\Big(0,\frac{1}{a}\Big), \tag{3.15}$$

and $\{e_i, \mathcal{I}e_j\}$ *are constant unit vectors orthogonal to each other.*

Proof. Let us denote by ∇, $\dot{\nabla}$ and by $\widetilde{\nabla}$ the Levi-Civita connections on $(\mathbb{S}^{2n+1}, g)$, $(\mathbb{S}^{2n+1}, g_0)$ and $(\mathbb{R}^{2n+2}, \langle , \rangle)$, respectively.

First we consider the case when γ is the biharmonic circle, that is $\kappa_1^2 = \frac{c+3}{4}$. Let $T = \gamma'$ be the unit tangent vector field (with respect to the metric g) along γ. Using the two Sasakian structures on $\mathbb{S}^{2n+1}$ we obtain $\dot{\nabla}_T T = \nabla_T T$ and $\dot{\nabla}_T E_2 = \nabla_T E_2$.

From the Gauss and Frenet equations we get

$$\widetilde{\nabla}_T T = \dot{\nabla}_T T - \langle T, T\rangle\gamma = \kappa_1 E_2 - \frac{1}{a}\gamma$$

and

$$\widetilde{\nabla}_T \widetilde{\nabla}_T T = (-\kappa_1^2 - \frac{1}{a})T = -\frac{2}{a}T.$$

Hence

$$a\gamma^{'''} + 2\gamma' = 0,$$

with the general solution

$$\gamma(s) = \cos\Big(\sqrt{\frac{2}{a}}s\Big)c_1 + \sin\Big(\sqrt{\frac{2}{a}}s\Big)c_2 + c_3,$$

where $\{c_i\}$ are constant vectors in $\mathbb{E}^{2n+2}$.
As γ verifies the following equations,

$$\langle\gamma,\gamma\rangle = 1,\ \langle\gamma',\gamma'\rangle = \frac{1}{a},\ \langle\gamma,\gamma'\rangle = 0,\ \langle\gamma',\gamma''\rangle = 0,\ \langle\gamma'',\gamma''\rangle = \frac{2}{a^2},\ \langle\gamma,\gamma''\rangle = -\frac{1}{a},$$

and in $s = 0$ we have $\gamma = c_1 + c_3$, $\gamma' = \sqrt{\frac{2}{a}}c_2$, $\gamma'' = -\frac{2}{a}c_1$, one obtains

$$c_{11} + 2c_{13} + c_{33} = 1,\ c_{22} = \frac{1}{2},\ c_{12} + c_{23} = 0,\ c_{12} = 0,\ c_{11} = \frac{1}{2}, c_{11} + c_{13} = \frac{1}{2},$$

where $c_{ij} = \langle c_i, c_j\rangle$. Consequently, we obtain that $\{c_i\}$ are orthogonal vectors in $\mathbb{E}^{2n+2}$ with $|c_1| = |c_2| = |c_3| = \frac{1}{\sqrt{2}}$.
Finally, using the facts that γ is a Legendre curve and $g(\nabla_{\gamma'}\gamma', \varphi\gamma') = 0$ one obtains easily that $\langle c_i, \mathcal{I}c_j\rangle = 0$ for any $i, j = 1, 2, 3$.

Now we assume that γ is a biharmonic helix, that is $\kappa_1^2 + \kappa_2^2 = \frac{c+3}{4}$, $\kappa_1^2 \in \Big(0, \frac{c+3}{4}\Big)$.
First we obtain $\dot{\nabla}_T T = \nabla_T T$, $\dot{\nabla}_T E_2 = \nabla_T E_2$ and $\dot{\nabla}_T E_3 = \nabla_T E_3$.
From the Gauss and Frenet equations we get

$$\widetilde{\nabla}_T T = \dot{\nabla}_T T - \langle T, T\rangle\gamma = \kappa_1 E_2 - \frac{1}{a}\gamma,$$

$$\widetilde{\nabla}_T \widetilde{\nabla}_T T = \kappa_1 \widetilde{\nabla}_T E_2 - \frac{1}{a}T = \kappa_1\Big(-\kappa_1 T + \kappa_2 E_3\Big) - \frac{1}{a}T = -\Big(\kappa_1^2 + \frac{1}{a}\Big)T + \kappa_1\kappa_2 E_3,$$

and

$$\begin{aligned}\widetilde{\nabla}_T \widetilde{\nabla}_T \widetilde{\nabla}_T T &= -\Big(\kappa_1^2 + \tfrac{1}{a}\Big)\widetilde{\nabla}_T T + \kappa_1\kappa_2\widetilde{\nabla}_T E_3 = -\Big(\kappa_1^2 + \tfrac{1}{a}\Big)\widetilde{\nabla}_T T - \kappa_1\kappa_2^2 E_2 \\ &= -\tfrac{2}{a}\gamma'' - \tfrac{1}{a}\kappa_2^2\gamma.\end{aligned}$$

Therefore

$$a\gamma^{iv} + 2\gamma'' + \kappa_2^2\gamma = 0,$$

and its general solution is

$$\gamma(s) = \cos(As)c_1 + \sin(As)c_2 + \cos(Bs)c_3 + \sin(Bs)c_4,$$

where A, B are given by (3.15) and $\{c_i\}$ are constant vectors in $\mathbb{E}^{2n+2}$.

The curve γ satisfies

$$\langle\gamma,\gamma\rangle=1,\ \langle\gamma',\gamma'\rangle=\frac{1}{a},\ \langle\gamma,\gamma'\rangle=0,\ \langle\gamma',\gamma''\rangle=0,\ \langle\gamma'',\gamma''\rangle=\frac{1+a\kappa_1^2}{a^2},$$
$$\langle\gamma,\gamma''\rangle=-\frac{1}{a},\ \langle\gamma',\gamma'''\rangle=-\frac{1+a\kappa_1^2}{a^2},\ \langle\gamma'',\gamma'''\rangle=0,\ \langle\gamma,\gamma'''\rangle=0,$$
$$\langle\gamma''',\gamma'''\rangle=\frac{3a\kappa_1^2+1}{a^3},$$

and in $s=0$ we have

$$\gamma=c_1+c_3,\quad \gamma'=Ac_2+Bc_4,\quad \gamma''=-A^2c_1-B^2c_3,\quad \gamma'''=-A^3c_2-B^3c_4.$$

Then, it follows

$$c_{11}+2c_{13}+c_{33}=1 \tag{3.16}$$

$$A^2c_{22}+2ABc_{24}+B^2c_{44}=\frac{1}{a} \tag{3.17}$$

$$Ac_{12}+Ac_{23}+Bc_{14}+Bc_{34}=0 \tag{3.18}$$

$$A^3c_{12}+AB^2c_{23}+A^2Bc_{14}+B^3c_{34}=0 \tag{3.19}$$

$$A^4c_{11}+2A^2B^2c_{13}+B^4c_{33}=\frac{1+a\kappa_1^2}{a^2} \tag{3.20}$$

$$A^2c_{11}+(A^2+B^2)c_{13}+B^2c_{33}=\frac{1}{a} \tag{3.21}$$

$$A^4c_{22}+(AB^3+A^3B)c_{24}+B^4c_{44}=\frac{1+a\kappa_1^2}{a^2} \tag{3.22}$$

$$A^5c_{12}+A^3B^2c_{23}+A^2B^3c_{14}+B^5c_{34}=0 \tag{3.23}$$

$$A^3c_{12}+A^3c_{23}+B^3c_{14}+B^3c_{34}=0 \tag{3.24}$$

$$A^6c_{22}+2A^3B^3c_{24}+B^6c_{44}=\frac{3a\kappa_1^2+1}{a^3} \tag{3.25}$$

where $c_{ij}=\langle c_i,c_j\rangle$.
The solution of the system given by (3.18), (3.19), (3.23) and (3.24) is

$$c_{12}=c_{23}=c_{14}=c_{34}=0.$$

From equations (3.16), (3.20) and (3.21) we get

$$c_{11}=\frac{1}{2},\quad c_{13}=0,\quad c_{33}=\frac{1}{2},$$

and, from (3.17), (3.22), (3.25),

$$c_{22}=\frac{1}{2},\quad c_{24}=0,\quad c_{44}=\frac{1}{2}.$$

We obtain that $\{c_i\}$ are orthogonal vectors in $\mathbb{E}^8$ with $|c_1|=|c_2|=|c_3|=|c_4|=\frac{1}{\sqrt{2}}$. Finally, since γ is a Legendre curve and $g(\nabla_{\gamma'}\gamma',\varphi\gamma')=0$, one obtains the conclusion. □

Just like for $\mathbb{S}^3$ (see [11]) one obtains for $\mathbb{S}^{2n+1}$ endowed with the modified Sasakian structure, with $0<a<1$ (that means $c>1$), the following result.

Theorem 3.14. *Let* $\gamma : I \to (\mathbb{S}^{2n+1},\varphi,\xi,\eta,g)$ *be a biharmonic Legendre curve parametrized by its arc length such that* $\nabla_{\gamma'}\gamma' = \sqrt{c-1}\varphi\gamma'$. *Then the equation of* γ *in the Euclidean space* $\mathbb{E}^{2n+2}$ *is*

$$\begin{aligned}\gamma(s) \quad = \quad & \sqrt{\tfrac{B}{A+B}}\cos(As)e_1 - \sqrt{\tfrac{B}{A+B}}\sin(As)\mathcal{I}e_1 \\ & + \sqrt{\tfrac{A}{A+B}}\cos(Bs)e_3 + \sqrt{\tfrac{A}{A+B}}\sin(Bs)\mathcal{I}e_3 \\ = \quad & \sqrt{\tfrac{B}{A+B}}\exp(-\mathrm{i}As)e_1 + \sqrt{\tfrac{A}{A+B}}\exp(\mathrm{i}Bs)e_3,\end{aligned}$$

where $\{e_i\}$ *are constant unit orthogonal vectors in* $\mathbb{E}^{2n+2}$ *and*

$$A = \sqrt{\frac{3-2a-2\sqrt{(a-1)(a-2)}}{a}}, \quad B = \sqrt{\frac{3-2a+2\sqrt{(a-1)(a-2)}}{a}}. \tag{3.26}$$

Remark 3.15. The ODE satisfied by proper-biharmonic Legendre curves in the $(2n+1)$-sphere, in the fourth case, may be also obtained but the computations are rather complicated.

4. Biharmonic submanifolds in Sasakian space forms

A method to obtain biharmonic submanifolds in a Sasakian space form is provided by the following Theorem.

Theorem 4.1. *Let* $(N^{2n+1},\varphi,\xi,\eta,g)$ *be a strictly regular Sasakian space form with constant* φ*-sectional curvature* c *and let* $\mathbf{i}: M \to N$ *be an* r*-dimensional integral submanifold of* N. *Consider*

$$F : \widetilde{M} = I \times M \to N, \quad F(t,p) = \phi_t(p) = \phi_p(t),$$

where $I = \mathbb{S}^1$ *or* $I = \mathbb{R}$ *and* $\{\phi_t\}_{t\in\mathbb{R}}$ *is the flow of the vector field* ξ. *Then* $F : (\widetilde{M}, \widetilde{g} = dt^2 + \mathbf{i}^*g) \to N$ *is a Riemannian immersion and it is proper-biharmonic if and only if* M *is a proper-biharmonic submanifold of* N.

Proof. From the definition of the flow of ξ we have

$$dF(t,p)\Big(\frac{\partial}{\partial t}\Big) = \frac{d}{ds}|_{s=t}\{\phi_p(s)\} = \dot{\phi}_p(t) = \xi(\phi_p(t)) = \xi(F(t,p)),$$

i.e. $\frac{\partial}{\partial t}$ is F-correlated to ξ and

$$\Big|dF(t,p)\Big(\frac{\partial}{\partial t}\Big)\Big| = |\xi(F(t,p))| = 1 = \Big|\frac{\partial}{\partial t}\Big|.$$

The vector $X_p \in T_pM$ can be identified to $(0,X_p) \in T_{(t',p)}(I\times M)$, for any $t' \in I$, and we have

$$dF_{(t,p)}(X_p) = (dF)_{(t,p)}(\dot{\gamma}(0)) = \frac{d}{ds}|_{s=0}\{\phi_t(\gamma(s))\} = (d\phi_t)_p(X_p).$$

Since ϕ_t is an isometry, we have $|dF_{(t,p)}(X_p)| = |(d\phi_t)_p(X_p)| = |X_p|$.
Moreover,

$$\begin{aligned} g\Big(dF_{(t,p)}\Big(\tfrac{\partial}{\partial t}\Big), dF_{(t,p)}(X_p)\Big) &= g(\xi(\phi_p(t)), (d\phi_t)_p(X_p)) \\ &= g((d\phi_t)_p(\xi_p), (d\phi_t)_p(X_p)) = g(\xi_p, X_p) \\ &= 0, \end{aligned}$$

so $F : (I \times M, dt^2 + \mathbf{i}^* g) \to N$ is a Riemannian immersion.

Let $F^{-1}(TN)$ be the pull-back bundle over $\widetilde{M}$ and ∇^F the pull-back connection determined by the Levi-Civita connection on N. We shall prove that

$$\tau(F)_{(t,p)} = (d\phi_t)_p(\tau(\mathbf{i})) \quad \text{and} \quad \tau_2(F)_{(t,p)} = (d\phi_t)_p(\tau_2(\mathbf{i})),$$

so, from the point of view of harmonicity and biharmonicity, $\widetilde{M}$ and M have the same behaviour.

We start with two remarks. Let $\sigma \in F^{-1}(TN)$ defined by $\sigma_{(t,p)} = (d\phi_t)_p(Z_p)$, where Z is a vector field along M, that is $Z_p \in T_pN$, $\forall p \in M$. We have

$$(\nabla^F_X \sigma)_{(t,p)} = (d\phi_t)_p(\nabla^N_X Z), \quad \forall X \in C(TM). \tag{4.1}$$

Then, if $\sigma \in F^{-1}(TN)$, it follows that $\varphi\sigma \in F^{-1}(TN)$, $(\varphi\sigma)_{(t,p)} = \varphi_{\phi_p(t)}(\sigma_{(t,p)})$, and

$$\nabla^F_{\frac{\partial}{\partial t}} \varphi\sigma = \varphi \nabla^F_{\frac{\partial}{\partial t}} \sigma. \tag{4.2}$$

Now, we consider $\{X_1, ..., X_r\}$ a local orthonormal frame field on U, where U is an open subset of M. The tension field of F is given by

$$\tau(F) = \nabla^F_{\frac{\partial}{\partial t}} dF\Big(\frac{\partial}{\partial t}\Big) - dF\Big(\nabla^{\widetilde{M}}_{\frac{\partial}{\partial t}} \frac{\partial}{\partial t}\Big) + \sum_{a=1}^{r} \{\nabla^F_{X_a} dF(X_a) - dF(\nabla^{\widetilde{M}}_{X_a} X_a)\}.$$

As

$$\nabla^F_{\frac{\partial}{\partial t}} dF\Big(\frac{\partial}{\partial t}\Big) = \nabla^N_\xi \xi = 0, \quad \nabla^{\widetilde{M}}_{\frac{\partial}{\partial t}} \frac{\partial}{\partial t} = \nabla^I_{\frac{\partial}{\partial t}} \frac{\partial}{\partial t} = 0,$$

$$(\nabla^F_{X_a} dF(X_a))_{(t,p)} = (d\phi_t)_p(\nabla^N_{X_a} X_a), \quad dF_{(t,p)}(\nabla^{\widetilde{M}}_{X_a} X_a) = (d\phi_t)_p(\nabla^M_{X_a} X_a)$$

we obtain

$$\tau(F)_{(t,p)} = (d\phi_t)_p(\tau(\mathbf{i})).$$

The next step is to prove that $\nabla^F_{\frac{\partial}{\partial t}} \tau(F) = -\varphi(\tau(F))$. Since $[\frac{\partial}{\partial t}, X_a] = 0$, $a = 1, ..., r$,

$$\nabla^F_{\frac{\partial}{\partial t}} dF(X_a) = \nabla^F_{X_a} dF\Big(\frac{\partial}{\partial t}\Big).$$

But

$$\begin{aligned} \Big(\nabla^F_{X_a} dF\Big(\tfrac{\partial}{\partial t}\Big)\Big)_{(t,p)} &= \nabla^N_{dF_{(t,p)}X_a} \xi = \nabla^N_{(d\phi_t)_p X_a} \xi = -\varphi((d\phi_t)_p(X_a)) \\ &= -(d\phi_t)_p(\varphi X_a), \end{aligned}$$

so

$$\Big(\nabla^F_{\frac{\partial}{\partial t}} dF(X_a)\Big)_{(t,p)} = -(d\phi_t)_p(\varphi X_a). \tag{4.3}$$

We note that

$$R^F\Big(\frac{\partial}{\partial t}, X_a\Big) dF(X_a) = \nabla^F_{\frac{\partial}{\partial t}} \nabla^F_{X_a} dF(X_a) - \nabla^F_{X_a} \nabla^F_{\frac{\partial}{\partial t}} dF(X_a)$$

and, on the other hand

$$\Big(R^F\Big(\frac{\partial}{\partial t}, X_a\Big) dF(X_a)\Big)_{(t,p)} = R^N_{\phi_t(p)}(\xi, (d\phi_t)_p(X_a))(d\phi_t)_p(X_a) = \xi.$$

Therefore

$$\nabla^F_{\frac{\partial}{\partial t}} \nabla^F_{X_a} dF(X_a) - \nabla^F_{X_a} \nabla^F_{\frac{\partial}{\partial t}} dF(X_a) = \xi \tag{4.4}$$

From (4.1) and (4.3) we get

$$\begin{aligned}\left(\nabla^F_{X_a}\nabla^F_{\frac{\partial}{\partial t}} dF(X_a)\right)_{(t,p)} &= -(d\phi_t)_p(\nabla^N_{X_a}\varphi X_a)\\ &= -(d\phi_t)_p(\xi+\varphi\nabla^N_{X_a}X_a)\end{aligned}\tag{4.5}$$

and from (4.3)

$$\begin{aligned}\left(\nabla^F_{\frac{\partial}{\partial t}} dF(\nabla^{\widetilde{M}}_{X_a}X_a)\right)_{(t,p)} &= \left(\nabla^F_{\frac{\partial}{\partial t}} dF(\nabla^{M}_{X_a}X_a)\right)_{(t,p)}\\ &= -(d\phi_t)_p(\varphi\nabla^M_{X_a}X_a).\end{aligned}\tag{4.6}$$

Replacing (4.5) and (4.6) in (4.4), we obtain

$$\begin{aligned}\xi &= \nabla^F_{\frac{\partial}{\partial t}}\nabla^F_{X_a} dF(X_a) - \nabla^F_{\frac{\partial}{\partial t}} dF(\nabla^{\widetilde{M}}_{X_a}X_a) + \nabla^F_{\frac{\partial}{\partial t}} dF(\nabla^{\widetilde{M}}_{X_a}X_a) - \nabla^F_{X_a}\nabla^F_{\frac{\partial}{\partial t}} dF(X_a)\\ &= \nabla^F_{\frac{\partial}{\partial t}}\nabla dF(X_a,X_a) - (d\phi_t)_p(\varphi\nabla^M_{X_a}X_a) + (d\phi_t)_p(\xi+\varphi\nabla^N_{X_a}X_a),\end{aligned}$$

so

$$\left(\nabla^F_{\frac{\partial}{\partial t}}\nabla^F_{X_a} dF(X_a)\right)_{(t,p)} = -\varphi(d\phi_t)_p(\nabla d\mathbf{i}(X_a,X_a)).$$

As $\nabla dF(X_a,X_a)=0$, we obtain

$$\nabla^F_{\frac{\partial}{\partial t}}\tau(F) = -\varphi(\tau(F)).$$

From (4.2) we have

$$\begin{aligned}\nabla^F_{\frac{\partial}{\partial t}}\nabla^F_{\frac{\partial}{\partial t}}\tau(F) &= -\nabla^F_{\frac{\partial}{\partial t}}\varphi(\tau(F)) = -\varphi\nabla^F_{\frac{\partial}{\partial t}}\tau(F) = \varphi^2\tau(F)\\ &= -\tau(F),\end{aligned}\tag{4.7}$$

and from (4.1)

$$(\nabla^F_{X_a}\nabla^F_{X_a}\tau(F))_{(t,p)} = (d\phi_t)_p(\nabla^N_{X_a}\nabla^N_{X_a}\tau(\mathbf{i})),\tag{4.8}$$

$$\left(\nabla^F_{\nabla^{\widetilde{M}}_{X_a}X_a}\tau(F)\right)_{(t,p)} = (d\phi_t)_p\left(\nabla^N_{\nabla^M_{X_a}X_a}\tau(\mathbf{i})\right).\tag{4.9}$$

From (4.7), (4.8) and (4.9) we obtain

$$-\,(\Delta^F\tau(F))_{(t,p)} = -\tau(F)_{(t,p)} - (d\phi_t)_p(\Delta^{\mathbf{i}}\tau(\mathbf{i})).\tag{4.10}$$

After a straightforward computation we get

$$\operatorname{trace} R^F(dF,\tau(F))dF = -\tau(F) + (d\phi_t)_p(\operatorname{trace} R^N_p(d\mathbf{i},\tau(\mathbf{i}))d\mathbf{i}).\tag{4.11}$$

Finally, from (4.10) and (4.11) we obtain

$$\tau_2(F)_{(t,p)} = (d\phi_t)_p(\tau_2(\mathbf{i})).$$

□

Remark 4.2. The previous result was expected because of the following simple remark. Assume that $(N^{2n+1},\varphi,\xi,\eta,g)$ is a compact strictly regular Sasakian space form with constant φ-sectional curvature c and let $G : M \to N$ be an arbitrary smooth map from a compact Riemannian manifold M. If F is biharmonic, then the map G is biharmonic, where $F : \widetilde{M} = \mathbb{S}^1\times M \to N$, $F(t,p) = \phi_t(G(p))$.

Indeed, an arbitrary variation $\{G_s\}_s$ of G induces a variation $\{F_s\}_s$ of F. We have that $\tau_{(p,t)}(F_s) = (d\phi_t)_{G_s(p)}(\tau_p(G_s))$ and, from the biharmonicity of F and the Fubini Theorem, we get

$$\begin{aligned} 0 &= \frac{d}{ds}|_{s=0}\{E_2(F_s)\} = \frac{1}{2}\frac{d}{ds}|_{s=0}\int_{\widetilde{M}} |\tau(F_s)|^2 \ v_{\widetilde{g}} = \frac{\pi}{2}\frac{d}{ds}|_{s=0}\int_M |\tau(G_s)|^2 \ v_g \\ &= \pi\frac{d}{ds}|_{s=0}\{E_2(G_s)\}. \end{aligned}$$

Next, consider the unit $(2n+1)$-dimensional sphere $\mathbb{S}^{2n+1}$ endowed with its canonical or modified Sasakian structure. The flow of ξ is $\phi_t(z) = \exp(-\mathrm{i}\frac{t}{a})z$, where

$$z = (z^1, ..., z^{n+1}) = (x^1, ..., x^{n+1}, y^1, ..., y^{n+1}).$$

From the above expression of the flow and from Theorems 3.14, 3.13, 3.12 and 4.1 we obtain explicit examples of proper-biharmonic submanifolds in $(\mathbb{S}^{2n+1}, \varphi, \xi, \eta, g)$, $a > 0$, of constant mean curvature. In particular, we reobtain a result in [1] and, for $n = 1$, the result in [11].

Proposition 4.3 ([1]). *Let $F : \widetilde{M} \to \mathbb{S}^5 \subset \mathbb{R}^6$ be a proper-biharmonic anti-invariant immersion. Then the position vector of $\widetilde{M}$ in $\mathbb{R}^6$ is*

$$F(t,u,v) = \frac{\exp(-\mathrm{i}t)}{\sqrt{2}}(\exp(\mathrm{i}u), \mathrm{i}\exp(-\mathrm{i}u)\sin(\sqrt{2}v), \mathrm{i}\exp(-\mathrm{i}u)\cos(\sqrt{2}v)).$$

Proof. It was proved in [17] that the proper-biharmonic integral surface of $(\mathbb{S}^5, \varphi_0, \xi_0, \eta_0, g_0)$ is given by

$$f(u,v) = \frac{1}{\sqrt{2}}(\exp(\mathrm{i}u), \mathrm{i}\exp(-\mathrm{i}u)\sin\sqrt{2}v, \mathrm{i}\exp(-\mathrm{i}u)\cos\sqrt{2}v).$$

Now, composing with the flow of ξ_0 we reobtain the result in [1]. □

Proposition 4.4 ([11]). *Let M be a surface in $(\mathbb{S}^{2n+1}, \varphi, \xi, \eta, g)$ with $a \in (0,1)$, with the position vector in the Euclidean space $\mathbb{E}^{2n+2}$ given by*

$$F(t,s) = \exp\Big(-\mathrm{i}\frac{t}{a}\Big)\Big(\sqrt{\frac{B}{A+B}}\exp(-\mathrm{i}As)e_1 + \sqrt{\frac{A}{A+B}}\exp(\mathrm{i}Bs)e_3\Big),$$

where $\{e_1, e_3\}$ is an orthonormal system of constant vectors in the Euclidean space $(\mathbb{R}^{2n+2}, \langle,\rangle)$, with e_3 orthogonal to $\mathcal{I}e_1$ and A, B are given by (3.26). Then M is a proper-biharmonic surface in $\mathbb{S}^{2n+1}(c)$.

5. PROPER-BIHARMONIC SUBMANIFOLDS OF $(\mathbb{S}^7, g_0)$

We consider the Euclidean space $\mathbb{E}^8$ with three complex structures,

$$\mathcal{I} = \begin{pmatrix} 0 & -I_4 \\ I_4 & 0 \end{pmatrix}, \quad \mathcal{J} = \begin{pmatrix} 0 & 0 & 0 & I_2 \\ 0 & 0 & -I_2 & 0 \\ 0 & I_2 & 0 & 0 \\ -I_2 & 0 & 0 & 0 \end{pmatrix}, \quad \mathcal{K} = -\mathcal{I}\mathcal{J},$$

where I_n denotes the $n \times n$ identity matrix. We define three vector fields on $\mathbb{S}^7$ by

$$\xi_1 = -\mathcal{I}z, \quad \xi_2 = -\mathcal{J}z, \quad \xi_3 = -\mathcal{K}z, \quad z \in \mathbb{S}^7,$$

and consider their dual 1-forms $\eta_1 = \eta_0$, η_2, η_3. Let φ_a defined by

$$\varphi_1 = \varphi_0 = s \circ \mathcal{I}, \quad \varphi_2 = s \circ \mathcal{J}, \quad \varphi_3 = s \circ \mathcal{K}.$$

Then $(\varphi_a, \xi_a, \eta_a, g_0)$, $a = 1, 2, 3$, determine a Sasakian 3-structure on $\mathbb{S}^7$.

Now, we shall indicate a method to construct proper-biharmonic submanifolds in $(\mathbb{S}^7, g_0)$. We consider $\gamma = \gamma(s)$ a proper-biharmonic curve in $(\mathbb{S}^7, g_0)$, parametrized by arc-length, which is a Legendre curve for two of the three contact structures (it was proved in [9] that there is no proper-biharmonic curve which is Legendre with respect to all three contact structures on $\mathbb{S}^7$). For example, assume that γ is a Legendre curve for η_1 and η_2. Composing with the flow of ξ_1 (or ξ_2) we obtain a biharmonic surface which is Legendre with respect to η_2 (or η_1). Then, composing with the flow of ξ_2 (or ξ_1) we get a biharmonic 3-dimensional submanifold of $(\mathbb{S}^7, g_0)$. Using this method, from Theorems 3.12 and 4.1, we obtain 4 classes of proper-biharmonic surfaces in $(\mathbb{S}^7, g_0)$ and 4 classes of proper-biharmonic 3-dimensional submanifolds of $(\mathbb{S}^7, g_0)$, all of constant mean curvature.
For example, from Theorems 3.12 and 4.1, composing first with the flow of ξ_1 and then with that of ξ_2, we get the explicit parametric equations of proper-biharmonic 3-dimensional submanifolds of $(\mathbb{S}^7, g_0)$.

Proposition 5.1. *Let M be a 3-dimensional submanifold in $\mathbb{S}^7$ such that its position vector field in $\mathbb{E}^8$ is either*

$$\begin{aligned}
\mathbf{x}_1 &= \mathbf{x}_1(u,t,s)\\
&= \tfrac{1}{\sqrt{2}}\Big(\cos(u)\cos(\sqrt{2}s)\cos(t)e_1 + \cos(u)\sin(\sqrt{2}s)\cos(t)e_2\\
&\quad + \cos(u)\cos(t)e_3 - \cos(u)\cos(\sqrt{2}s)\sin(t)\mathcal{I}e_1\\
&\quad - \cos(u)\sin(\sqrt{2}s)\sin(t)\mathcal{I}e_2 - \cos(u)\sin(t)\mathcal{I}e_3\\
&\quad - \sin(u)\cos(\sqrt{2}s)\cos(t)\mathcal{J}e_1 - \sin(u)\sin(\sqrt{2}s)\cos(t)\mathcal{J}e_2\\
&\quad - \sin(u)\cos(t)\mathcal{J}e_3 - \sin(u)\cos(\sqrt{2}s)\sin(t)\mathcal{K}e_1\\
&\quad - \sin(u)\sin(\sqrt{2}s)\sin(t)\mathcal{K}e_2 - \sin(u)\sin(t)\mathcal{K}e_3\Big),
\end{aligned}$$

where $\{e_i, \mathcal{I}e_j\}$, $\{e_i, \mathcal{J}e_j\}$ are systems of constant orthonormal vectors in $\mathbb{E}^8$, or

$$\begin{aligned}
\mathbf{x}_2 &= \mathbf{x}_2(u,t,s)\\
&= \tfrac{1}{\sqrt{2}}\Big(\cos(u)\cos(As)\cos(t)e_1 + \cos(u)\sin(As)\cos(t)e_2\\
&\quad + \cos(u)\cos(Bs)\cos(t)e_3 + \cos(u)\sin(Bs)\cos(t)e_4\\
&\quad - \cos(u)\cos(As)\sin(t)\mathcal{I}e_1 - \cos(u)\sin(As)\sin(t)\mathcal{I}e_2\\
&\quad - \cos(u)\cos(Bs)\sin(t)\mathcal{I}e_3 - \cos(u)\sin(Bs)\sin(t)\mathcal{I}e_4\\
&\quad - \sin(u)\cos(As)\cos(t)\mathcal{J}e_1 - \sin(u)\sin(As)\cos(t)\mathcal{J}e_2\\
&\quad - \sin(u)\cos(Bs)\cos(t)\mathcal{J}e_3 - \sin(u)\sin(Bs)\cos(t)\mathcal{J}e_4\\
&\quad - \sin(u)\cos(As)\sin(t)\mathcal{K}e_1 - \sin(u)\sin(As)\sin(t)\mathcal{K}e_2\\
&\quad - \sin(u)\cos(Bs)\sin(t)\mathcal{K}e_3 - \sin(u)\sin(Bs)\sin(t)\mathcal{K}e_4\Big),
\end{aligned}$$

where

$$A=\sqrt{1+\kappa_1},\quad B=\sqrt{1-\kappa_1},\quad \kappa_1=\text{constant}\in(0,1),$$

and $\{e_i\}$ *are constant orthonormal vectors in* $\mathbb{E}^8$ *such that*

$$\langle e_1,\mathcal{I}e_3\rangle=\langle e_1,\mathcal{I}e_4\rangle=\langle e_2,\mathcal{I}e_3\rangle=\langle e_2,\mathcal{I}e_4\rangle=0,$$

$$\langle e_1,\mathcal{J}e_3\rangle=\langle e_1,\mathcal{J}e_4\rangle=\langle e_2,\mathcal{J}e_3\rangle=\langle e_2,\mathcal{J}e_4\rangle=0,$$

and

$$A\langle e_1,\mathcal{I}e_2\rangle+B\langle e_3,\mathcal{I}e_4\rangle=A\langle e_1,\mathcal{J}e_2\rangle+B\langle e_3,\mathcal{J}e_4\rangle=0.$$

Then M *is a proper-biharmonic submanifold of* $(\mathbb{S}^7,g_0)$.

Proof. As the flows of ξ_1 and ξ_2 are given by

$$\phi^1_t(z)=(\cos t)z-(\sin t)\mathcal{I}z,\qquad \phi^2_t(z)=(\cos t)z-(\sin t)\mathcal{J}z,$$

the Proposition follows by a straightforward computation.

□

Remark 5.2. Note that there exist vectors $\{e_i\}$ which satisfy the hypotheses of the previous Proposition. For example the first three or four vectors, respectively, from the canonical basis of $\mathbb{E}^8$ satisfy the required properties.

REFERENCES

[1] K. Arslan, R. Ezentas, C. Murathan, T. Sasahara. *3-dimensional biharmonic anti-invariant submanifolds in Sasakian space forms*, Beiträge Algebra Geom., to appear.

[2] A. Arvanitoyeorgos, F. Defever, G. Kaimakamis, V.J. Papantoniou. *Biharmonic Lorentz hypersurfaces in* E^4_1, Pacific J. Math., 229(2007), 293–305.

[3] C. Baikoussis, D.E. Blair. *On the geometry of 7-sphere*, Results in Math., 27(1995), 5–16.

[4] C. Baikoussis, D.E. Blair, T. Koufogiorgos. *Integral submanifolds of Sasakian space forms* $\bar{M}^7$, Results in Math., 27(1995), 207–226.

[5] A. Balmuş, S. Montaldo, C. Oniciuc. *Classification results for biharmonic submanifolds in spheres*, Israel J. Math., to appear.

[6] D.E. Blair. *Riemannian Geometry of Contact and Symplectic Manifolds*, Birkhäuser Boston, Progress in Mathematics, Volume 203, 2002.

[7] B.Y. Chen. *A report on submanifolds of finite type*, Soochow J. Math., 22(1996), 117–337.

[8] J. Eells, J.H. Sampson. *Harmonic mappings of Riemannian manifolds*, Amer. J. Math., 86(1964), 109–160.

[9] D. Fetcu. *Integral submanifolds in 3-Sasakian manifolds whose mean curvature vector fields are eigenvectors of the Laplace operator*, preprint.

[10] D. Fetcu. *Biharmonic Legendre curves in Sasakian space forms*, preprint.

[11] D. Fetcu, C. Oniciuc. *Explicit formulas for biharmonic submanifolds in non-Euclidean 3-spheres*, Abh. Math. Semin. Univ. Hamb., to appear.

[12] T. Ichiyama, J. Inoguchi, H. Urakawa. *Bi-harmonic maps and bi-Yang-Mills fields*, preprint.

[13] J. Inoguchi. *Submanifolds with harmonic mean curvature in contact 3-manifolds*, Colloq. Math., 100(2004), 163–179.

[14] G.Y. Jiang. *2-harmonic maps and their first and second variational formulas*, Chinese Ann. Math. Ser. A7(4)(1986), 389–402.

[15] S. Montaldo, C. Oniciuc. *A short survey on biharmonic maps between Riemannian manifolds*, Rev. Un. Mat. Argentina, 47(2)(2006), 1–22.

[16] Y-L. Ou, Z-P. Wang. *Biharmonic maps into Sol and Nil spaces*, `arXiv:math.DG/0612329v1`.

[17] T. Sasahara. *Legendre surfaces in Sasakian space forms whose mean curvature vectors are eigenvectors*, Publ. Math. Debrecen, 67(3–4)(2005), 285–303.

[18] The Bibliography of Biharmonic Maps. `http://beltrami.sc.unica.it/ biharmonic/`

[19] S. Tanno. *The topology of contact Riemannian manifolds*, Ill. J. Math., 12(1968), 700–717.

[20] W. Zhang. *New examples of biharmonic submanifolds in* $\mathbb{C}P^n$ *and* $\mathbb{S}^{2n+1}$, `arXiv:math.DG/07053961v1`.

Department of Mathematics, "Gh. Asachi" Technical University of Iasi, Bd. Carol I no. 11, 700506 Iasi, Romania
E-mail address: `dfetcu@math.tuiasi.ro`

Faculty of Mathematics, "Al.I. Cuza" University of Iasi, Bd. Carol I no. 11, 700506 Iasi, Romania
E-mail address: `oniciucc@uaic.ro`